arXiv:quant-ph/9708005v1 3 Aug 1997

Quantum Database Searching by a Single Query

Dong Pyo Chi [*][†] and Jinsoo Kim [*][‡]

15 July 1997

Abstract

In this paper we give a quantum mechanical algorithm that can search a database by a single query, when the number of solutions is more than a quarter. It utilizes modified Grover operator of arbitrary phase.

1 Introduction

For $N \in \mathbb{N}$, let $\mathbb{Z}_N = \{0, 1, \ldots, N-1\}$ denote the additive cyclic group of order N and consider an arbitrary function $F : \mathbb{Z}_N \to \mathbb{Z}_2$, The database searching problem is to find some $i \in \mathbb{Z}_N$ such that $F(i) = 1$ under the assumption that such an i exists. We assume that the structure of F is unknown so that it is not possible to obtain a knowledge about F without evaluating it on $\mathbb{Z}_N$.

Let $t = |\{i \in \mathbb{Z}_N : F(i) = 1\}|$. There is a quantum mechanical algorithm to solve this problem in expected time of order $O(\sqrt{N/t})$, which is optimal up to a multiplicative constant [4, 2, 1]. Especially when $t = N/4$ is known, the original Grover algorithm in [4] can search a solution only by a single query [2]. It uses the π-phase, i.e., marking the states by multiplying $e^{\pi\imath} = -1$. When $t = N/2$, by changing this phase to $\pi/2$, that is, by marking the states by multiplying $e^{\pi\imath/2} = \imath$ and modifying the corresponding diffusion transform according to this phase, the solution can be found with certainty after a single iteration [3].

In this paper, we generalize Grover algorithm for arbitrary phase. When $t \geq N/4$, we give generalized conditional phase and diffusion transform

---

[*] Department of Mathematics, Seoul National University, Seoul 151-742, Korea.

[†] E-mail address: `dpchi@math.snu.ac.kr`

[‡] E-mail address: `jskim@math.snu.ac.kr`

depending on t, and then formulate a quantum mechanical algorithm that solves the database searching problem in a single query.

2 Grover Operator of Arbitrary Phase

Let $\mathcal{B}_N = \{|a\rangle\}_{a\in\mathbb{Z}_2^n} = \{|e_1\rangle, |e_2\rangle, \ldots, |e_N\rangle\}$ be the standard basis of an n-qubit quantum register with $N = 2^n$ and $\mathcal{H}_N$ be the corresponding Hilbert space, which represents the state vectors of a quantum system. Let $\mathcal{H}_m$ be an m-dimensional subspace of $\mathcal{H}_N$ spanned by a basis $\mathcal{B}_m = \{|e_{i_1}\rangle, |e_{i_2}\rangle, \ldots, |e_{i_m}\rangle\}$. Let l be a positive integer such that $1 \le l \le m$.

For $\gamma \in \mathbb{R}$, the *conditional γ-phase transform* on a subspace $\mathcal{H}_m$, $\mathbf{S}_{F,\gamma}^{\mathcal{H}_m} : \mathcal{H}_m \to \mathcal{H}_m$ is defined by

$$\mathbf{S}_{F,\gamma}^{\mathcal{H}_m}|e_{i_k}\rangle = (e^{\imath\gamma})^{F(e_{i_k})}|e_{i_k}\rangle\,,$$

for $k = 1, 2, \ldots, m$, where $\imath = \sqrt{-1}$. Let $\mathbf{S}_{l,\gamma}^{\mathcal{H}_m}$ denote $\mathbf{S}_{F_l,\gamma}^{\mathcal{H}_m}$, where $F_l(e_{i_k}) = \delta_{i_k i_l}$.

Let $\mathbf{W}_l^{\mathcal{H}_m}$ be any unitary transformation on $\mathcal{H}_m$ satisfying

$$\mathbf{W}_l^{\mathcal{H}_m}|e_{i_l}\rangle = \frac{1}{\sqrt{m}}\sum_{k=1}^{m}|e_{i_k}\rangle\,.$$

For example, when m is a power of 2, in a suitably arranged basis we may set $\mathbf{W}_1^{\mathcal{H}_m}$ to be the Walsh-Hadamard transform. When m is not a power of 2, the approximate Fourier transform in [5] can be used.

For $\beta \in \mathbb{R}$, the *β-phase diffusion transform* on a subspace $\mathcal{H}_m$, $\mathbf{D}_\beta^{\mathcal{H}_m} : \mathcal{H}_N \to \mathcal{H}_N$ is defined by

$$\mathbf{D}_{\beta,ij}^{\mathcal{H}_m} = \langle e_i|\mathbf{D}_\beta^{\mathcal{H}_m}|e_j\rangle = \begin{cases} \frac{e^{\imath\beta}-1}{m} & \text{when } |e_i\rangle, |e_j\rangle \in \mathcal{B}_m \text{ and } i \neq j, \\ 1 + \frac{e^{\imath\beta}-1}{m} & \text{when } |e_i\rangle, |e_j\rangle \in \mathcal{B}_m \text{ and } i = j, \\ \delta_{ij} & \text{otherwise}\,. \end{cases}$$

If we rearrange the basis $\mathcal{B}_N$ so that $\mathcal{B}_m = \{|e_1\rangle, |e_2\rangle, \ldots, |e_m\rangle\}$ and represent $\mathbf{D}_\beta^{\mathcal{H}_m}$ by its matrix $\mathbf{D}_\beta^{\mathcal{H}_m} = (\mathbf{D}_{\beta,ij}^{\mathcal{H}_m})$, then we have

$$\begin{aligned} \mathbf{D}_\beta^{\mathcal{H}_m} &= \begin{bmatrix} \mathbf{W}_l^{\mathcal{H}_m}\mathbf{S}_{l,\beta}^{\mathcal{H}_m}\mathbf{W}_l^{\mathcal{H}_m\dagger} & 0 \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} I + (e^{\imath\beta} - 1)P^{\mathcal{H}_m} & 0 \\ 0 & I \end{bmatrix}, \end{aligned}$$

where $P^{\mathcal{H}_m} = (P^{\mathcal{H}_m}_{ij})$ is a projection matrix in $\mathcal{H}_m$ with $P^{\mathcal{H}_m}_{ij} = \frac{1}{m}$. Note that $\mathbf{S}^{\mathcal{H}_m}_{F,\gamma}$ and $\mathbf{D}^{\mathcal{H}_m}_{\beta}$ are unitary.

Let $\mathbf{G}^{\mathcal{H}_m}_{F,\beta,\gamma} : \mathcal{H}_N \to \mathcal{H}_N$ be the *Grover operator* of (β, γ)-phase in $\mathcal{H}_m$ defined by

$$\mathbf{G}^{\mathcal{H}_m}_{F,\beta,\gamma} = \mathbf{D}^{\mathcal{H}_m}_{\beta}\mathbf{S}^{\mathcal{H}_m}_{F,\gamma} .$$

When $\beta = \gamma$ set $\mathbf{G}^{\mathcal{H}_m}_{F,\gamma} = \mathbf{G}^{\mathcal{H}_m}_{F,\gamma,\gamma}$. For simplicity, we shall assume that $m = N$ and omit the superscript $\mathcal{H}_m$.

Let

$$\begin{aligned} A &= \{e_j \in \mathbb{Z}_2^n | F(e_j) = 1\} , \\ B &= \{e_j \in \mathbb{Z}_2^n | F(e_j) = 0\} , \end{aligned}$$

and let $t = |A|$. For $k, l \in \mathbb{C}$ such that $t|k|^2 + (N - t)|l|^2 = 1$, define

$$|\psi(k, l)\rangle = \sum_{e_j \in A} k|e_j\rangle + \sum_{e_j \in B} l|e_j\rangle .$$

Lemma 1 *For $\beta, \gamma \in \mathbb{R}$, let*

$$|\psi(k_j, l_j)\rangle = \mathbf{G}^{j}_{F,\beta,\gamma}|\psi(k_0, l_0)\rangle .$$

Then after applying $j + 1$ Grover operator of (β, γ)-phase, we have

$$\begin{cases} k_{j+1} = \dfrac{(e^{\imath\beta} - 1)t + N}{N} e^{\imath\gamma} k_j + \dfrac{(e^{\imath\beta} - 1)(N - t)}{N} l_j , \\ l_{j+1} = \dfrac{(e^{\imath\beta} - 1)t}{N} e^{\imath\gamma} k_j + \dfrac{(e^{\imath\beta} - 1)(N - t) + N}{N} l_j . \end{cases} \tag{2.1}$$

Theorem 1 *Assume that $k_0 = l_0$ and $\beta, \gamma \in [0, 2\pi]$. Then $l_1 = 0$ if and only if $\frac{N}{4} \le t \le N$ and $\beta = \gamma = \cos^{-1}\left(1 - \frac{N}{2t}\right)$. In this case, $k_1 = (e^{\imath\gamma} - 1)k_0$.*

Proof. By (2.1), we get

$$\begin{cases} k_1 = \left[(e^{\imath\beta} - 1)(e^{\imath\gamma} - 1)\dfrac{t}{N} + e^{\imath\gamma} + e^{\imath\beta} - 1\right] k_0 , \\ l_1 = \left[(e^{\imath\beta} - 1)(e^{\imath\gamma} - 1)\dfrac{t}{N} + e^{\imath\beta}\right] l_0 . \end{cases} \tag{2.2}$$

By considering the imaginary part of $e^{-\imath\beta}\frac{l_1}{l_0}$, the equation

$$\frac{t}{N}\left\{(1-\cos\beta)\sin\gamma+(\cos\gamma-1)\sin\beta\right\}=0$$

is equivalent to

$$\frac{1-\cos\beta}{\sin\beta}=\frac{1-\cos\gamma}{\sin\gamma}\,. \tag{2.3}$$

Considering the real part of $e^{-\imath\beta}\frac{l_1}{l_0}$, from (2.3) it follows that the equation

$$\frac{t}{N}\left\{(1-\cos\beta)(1-\cos\gamma)+\sin\beta\sin\gamma\right\}-1=0$$

is equivalent to

$$\cos\beta=\cos\gamma=1-\frac{N}{2t}\,.$$

Again by (2.3), we get $\beta=\gamma$ and $t\geq\frac{N}{4}$. Furthermore, by (2.2) we obtain $k_1=(e^{\imath\gamma}-1)k_0$. This completes the proof. $\square$

For the case of π-phase in [4], $-\mathbf{D}_\pi=-I+2P$ is an inversion about average operation and we have

$$-\mathbf{D}_\pi|\psi(k,l)\rangle=\left|\psi(-\tfrac{N-2t}{N}k+\tfrac{2(N-t)}{N}l,\tfrac{N-2t}{N}l+\tfrac{2t}{N}k)\right\rangle.$$

In this case, there is an explicit closed-form formula for k_j and l_j;

$$\begin{cases} k_j=(-1)^j\dfrac{1}{\sqrt{t}}\sin\left((2j+1)\theta\right)\,, \\ l_j=(-1)^j\dfrac{1}{\sqrt{N-t}}\cos\left((2j+1)\theta\right)\,, \end{cases}$$

for $j=0,1,\ldots,$ where the angle θ is defined so that $\sin^2\theta=\frac{t}{N}$ [2]. Especially when $t=N/4$, we have $l_1=0$.

Grover operator of $\frac{\pi}{2}$-phase was used in [3]. Since

$$\mathbf{D}_{\frac{\pi}{2}}|\psi(k,l)\rangle=\left|\psi(\tfrac{(\imath-1)t+N}{N}k+\tfrac{(\imath-1)(N-t)}{N}l,\tfrac{(\imath-1)(N-t)+N}{N}l+\tfrac{(\imath-1)t}{N}k)\right\rangle,$$

when $t=\frac{N}{2}$, we have

$$\mathbf{G}_{F,\frac{\pi}{2}}|\psi(k,k)\rangle=|\psi((\imath-1)k,0)\rangle\,.$$

By Theorem 1, When $t\in[N/4,N]$, we have

$$\mathbf{G}_{F,\gamma}|\psi(k_0,k_0)\rangle=|\psi((e^{\imath\gamma}-1)k_0,0)\rangle\,,$$

where the phase γ is defined by $\gamma=\cos^{-1}\left(1-\frac{N}{2t}\right)$.

References

[1] C. H. Bennett, E. Bernstein, G. Brassard and U. Vazirani, *Strengths and weaknesses of quantum computing*, Los Alamos e-print quant-ph/9701001, 1997.

[2] M. Boyer, G. Brassard, P. Høyer and A. Tapp, *Tight bounds on quantum searching*, Proceedings of the Fourth Workshop on Physics and Computation, New England Complex Systems Institute, 1996, pp. 36–43; Los Alamos e-print quant-ph/9605034, 1996.

[3] G. Brassard and P. Høyer, *An exact quantum polynomial-time algorithm for Simon's problem*, Proceedings of the Fifth Israeli Symposium on Theory of Computing and Systems, 1997, (to appear); Los Alamos e-print quant-ph/9704027, 1997.

[4] L. K. Grover, *Quantum mechanics helps in searching for a needle in a haystack*, Phys. Rev. Lett., (to appear); Los Alamos e-print quant-ph/9605043, 1996.

[5] A. Yu. Kitaev, *Quantum measurements and the abelian stabilizer problem*, Los Alamos e-print quant-ph/9511026, 1995.